

Symptomwirkung	THC	CBD
Anticonvulsiv	+	++
Muskelrelaxierend	++	+
Antinozizeptiv / analgetisch	++	+
Sedierend	+	+
Angstlösend	+	++
Antidepressiv	+	-
Antiinflammatorisch	+	++
Spastikminderung	++	+
Appetitsteigerung	+	-
Antiemetisch	++	+
Antipsychotisch	-	++
Abnahme des Tränenflusses	+	-
Senkung des Augeninnendrucks	+	-
Antidyskinetisch	+	+
Tremorreduktion	+	-
Bronchospasmolytisch	+	-
Verminderte Speichel-Produktion	+	-
Hemmung der Magensäure-Produktion	++	+
Entspannung	+	++
Antiagitativ	+	-

modifiziert nach Russo et al. 2006

© Medical Cannabis Training 2019, T. Vatterrodt

Quellen:

- Brian Brown; PupilSize After Use of Marijuana and Alcohol: American Journal of Ophthalmology April 1977
- Franjo Grotenhermen; Clinical Pharmacodynamics of Cannabinoids: Journal of Cannabis Therapeutics, Vol. 4(1) 2004
- Thomas F. Denson, Mitchell Earleywine; Decreased depression in marijuana users, AddictiveBehaviors31 (2006) 738-742
- Hanan Abramovici Ph.D.; INFORMATION FOR HEALTH CARE PROFESSIONALS Cannabis (marihuana, marijuana) and the cannabinoids
- C. B. Carroll, P. G. Bain, L. Teare, et al.; Cannabis for dyskinesia in Parkinson disease: A randomized double-blind crossover study; Neurology2004;63;1245
- Clifford DB; Tetrahydrocannabinol for tremorin multiple sclerosis. Ann Neurol 13:669-67 1, 1983
- Paul Consroe, Reuven Sandyk & Stuart R. Snider(1986) Open label evaluation of cannabidiol in dystonic movement disorders, International Journal of Neuroscience, 30:4, 277-282
- Landschaft et al. (2017). Medical Grade Cannabis Clinical Guide. The Israeli Medical Cannabis Agency (IMCA), Ministry of Health. Draft edition 1.1 - January 2017
- Russo, Guy, Medical Hypotheses (2006) 66, 234-246